

PROTOCOL

Petaka coating with Matrigel

1 Thaw frozen Matrigel leaving it at 4°C overnight.



2 Prepare 10 Petakas ready to be used with a couple of stands.




3 Prepare a sterile 30 mL syringe, and Celartia's tips.




4 Prepare 20 ml of Diluted Matrigel (1 mg/ml) in serum-free cold cell culture media (RPMI1640, EMEM, DMEM, or others).




5 Introduce the 20 mL of diluted Matrigel into a Petaka.




6 Squeeze the Petaka until the media reaches the upper limit of the Petaka chamber, and hold for 10 seconds, then release.




7 Leave Petaka in a vertical position for 5 minutes.




8 Slowly withdraw the Matrigel solution from Petaka.



- 9 Put the Petaka in a completely horizontal position whilst all Petakas are being prepared



10




Introduce the residual Matrigel solution into the next Petaka.

11

Squeeze the new petaka until the media reaches the upper limit of the petaka chamber, and hold 10 seconds.


Repeat the steps 6 to 10 for the remaining Petakas



Notice that each Petaka treated will reduce the volume of Matrigel solution by around 2 to 3 mL. Therefore it is possible to treat efficiently about 5 Petakas covering about 80% of the available culture surface of the Petaka's chamber, which will lead to a residual 5 to 10 mL of Matrigel solution that could be saved at -80°C , until it is reused.

12

Incubate Petakas, in a horizontal position, at 37°C for 6 hours before seeding cells



Phase contrast microscopy of cells inside Petaka cultured on Matrigel

